

Some Triple Difference Rough Cesàro and Lacunary Statistical Sequence Spaces

Ayhan Esi and Nagarajan Subramanian

ABSTRACT: We generalized the concepts in probability of rough Cesàro and lacunary statistical by introducing the difference operator Δ_γ^α of fractional order, where α is a proper fraction and $\gamma = (\gamma_{mnk})$ is any fixed sequence of nonzero real or complex numbers. We study some properties of this operator involving lacunary sequence θ and arbitrary sequence $p = (p_{rst})$ of strictly positive real numbers and investigate the topological structures of related with triple difference sequence spaces.

The main focus of the present paper is to generalized rough Cesàro and lacunary statistical of triple difference sequence spaces and investigate their topological structures as well as some inclusion concerning the operator Δ_γ^α .

AMS Subject Classification: 40F05, 40J05, 40G05.

Keywords and Phrases: Analytic sequence; Musielak-Orlicz function; Triple sequences; Chi sequence; Cesàro summable; Lacunary statistical convergence.

Introduction

A triple sequence (real or complex) can be defined as a function $x : \mathbb{N} \times \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{R} (\mathbb{C})$, where \mathbb{N}, \mathbb{R} and \mathbb{C} denote the set of natural numbers, real numbers and complex numbers respectively. The different types of notions of triple sequence was introduced and investigated at the initial by Sahiner et al. [10, 11], Esi et al. [1-3], Dutta et al. [4], Subramanian et al. [12-15], Debnath et al. [5] and many others.

A triple sequence $x = (x_{mnk})$ is said to be triple analytic if

$$\sup_{m,n,k} |x_{mnk}|^{\frac{1}{m+n+k}} < \infty.$$

The space of all triple analytic sequences are usually denoted by Λ^3 . A triple sequence $x = (x_{mnk})$ is called triple gai sequence if

$$((m+n+k)!|x_{mnk}|)^{\frac{1}{m+n+k}} \rightarrow 0 \text{ as } m, n, k \rightarrow \infty.$$

The notion of difference sequence spaces (for single sequences) was introduced by Kizmaz [6] as follows

$$Z(\Delta) = \{x = (x_k) \in w : (\Delta x_k) \in Z\}$$

for $Z = c, c_0$ and ℓ_∞ , where $\Delta x_k = x_k - x_{k+1}$ for all $k \in \mathbb{N}$.

The difference triple sequence space was introduced by Debnath et al. (see [5]) and is defined as

$$\begin{aligned} \Delta x_{mnk} &= x_{mnk} - x_{m,n+1,k} - x_{m,n,k+1} + x_{m,n+1,k+1} \\ &\quad - x_{m+1,n,k} + x_{m+1,n+1,k} + x_{m+1,n,k+1} - x_{m+1,n+1,k+1} \end{aligned}$$

and $\Delta^0 x_{mnk} = \langle x_{mnk} \rangle$.

1. Some New Difference Triple Sequence Spaces with Fractional Order

Let $\Gamma(\alpha)$ denote the Euler gamma function of a real number α . Using the definition $\Gamma(\alpha)$ with $\alpha \notin \{0, -1, -2, -3, \dots\}$ can be expressed as an improper integral as follows: $\Gamma(\alpha) = \int_0^\infty e^{-x} x^{\alpha-1} dx$, where α is a positive proper fraction. We have defined the generalized fractional triple sequence spaces of difference operator

$$\Delta_\gamma^\alpha(x_{mnk}) = \sum_{u=0}^\infty \sum_{v=0}^\infty \sum_{w=0}^\infty \frac{(-1)^{u+v+w} \Gamma(\alpha+1)}{(u+v+w)! \Gamma(\alpha - (u+v+w) + 1)} x_{m+u, n+v, k+w}. \quad (1.1)$$

In particular, we have

$$\begin{aligned} (i) \quad \Delta^{\frac{1}{2}}(x_{mnk}) &= x_{mnk} - \frac{1}{16} x_{m+1, n+1, k+1} - \dots; \\ (ii) \quad \Delta^{-\frac{1}{2}}(x_{mnk}) &= x_{mnk} + \frac{5}{16} x_{m+1, n+1, k+1} + \dots; \\ (iii) \quad \Delta^{\frac{2}{3}}(x_{mnk}) &= x_{mnk} - \frac{4}{81} x_{m+1, n+1, k+1} - \dots. \end{aligned}$$

Now we determine the new classes of triple difference sequence spaces $\Delta_\gamma^\alpha(x)$ as follows:

$$\Delta_\gamma^\alpha(x) = \{x : (x_{mnk}) \in w^3 : (\Delta_\gamma^\alpha x) \in X\}, \quad (1.2)$$

where

$$\Delta_\gamma^\alpha(x_{mnk}) = \sum_{u=0}^\infty \sum_{v=0}^\infty \sum_{w=0}^\infty \frac{(-1)^{u+v+w} \Gamma(\alpha+1)}{(u+v+w)! \Gamma(\alpha - (u+v+w) + 1)} x_{m+u, n+v, k+w}$$

and

$$\begin{aligned} X \in \chi_f^{3\Delta}(x) &= \chi_f^3(\Delta_\gamma^\alpha x_{mnk}) = \mu_{mnk}(\Delta_\gamma^\alpha x) \\ &= \left[f_{mnk} \left(((m+n+k)! |\Delta_\gamma^\alpha|)^{\frac{1}{m+n+k}}, \bar{0} \right) \right]. \end{aligned}$$

Proposition 1.1.

- (i) For a proper fraction α $\Delta^\alpha : W \times W \times W \rightarrow W \times W \times W$ defined by equation of (2.1) is a linear operator;
- (ii) For $\alpha, \beta > 0$, $\Delta^\alpha (\Delta^\beta (x_{mnk})) = \Delta^{\alpha+\beta} (x_{mnk})$ and $\Delta^\alpha (\Delta^{-\alpha} (x_{mnk})) = x_{mnk}$.

Proof: Omitted.

Proposition 1.2. For a proper fraction α and f be an Musielak-Orlicz function, if $\chi_f^3(x)$ is a linear space, then $\chi_f^{3\Delta^\alpha}(x)$ is also a linear space.

Proof: Omitted.

2. Definitions and Preliminaries

Throughout the article $w^3, \chi^3(\Delta), \Lambda^3(\Delta)$ denote the spaces of all, triple gai difference sequence spaces and triple analytic difference sequence spaces respectively. Subramanian et al. (see [12]) introduced by a triple entire sequence spaces, triple analytic sequences spaces and triple gai sequence spaces. The triple sequence spaces of $\chi^3(\Delta), \Lambda^3(\Delta)$ are defined as follows:

$$\begin{aligned} \chi^3(\Delta) &= \left\{ x \in w^3 : ((m+n+k)! |\Delta x_{mnk}|)^{1/m+n+k} \rightarrow 0 \text{ as } m, n, k \rightarrow \infty \right\}, \\ \Lambda^3(\Delta) &= \left\{ x \in w^3 : \sup_{m,n,k} |\Delta x_{mnk}|^{1/m+n+k} < \infty \right\}. \end{aligned}$$

Definition 2.1. An Orlicz function ([see [7]) is a function $M : [0, \infty) \rightarrow [0, \infty)$ which is continuous, non-decreasing and convex with $M(0) = 0, M(x) > 0$, for $x > 0$ and $M(x) \rightarrow \infty$ as $x \rightarrow \infty$. If convexity of Orlicz function M is replaced by $M(x+y) \leq M(x) + M(y)$, then this function is called modulus function.

Lindenstrauss and Tzafriri ([8]) used the idea of Orlicz function to construct Orlicz sequence space.

A sequence $g = (g_{mn})$ defined by

$$g_{mn}(v) = \sup \{ |v| u - (f_{mnk})(u) : u \geq 0 \}, \quad m, n, k = 1, 2, \dots$$

is called the complementary function of a Musielak-Orlicz function f . For a given Musielak-Orlicz function f , (see [9]) the Musielak-Orlicz sequence space t_f is defined as follows

$$t_f = \left\{ x \in w^3 : I_f(|x_{mnk}|)^{1/m+n+k} \rightarrow 0 \text{ as } m, n, k \rightarrow \infty \right\},$$

where I_f is a convex modular defined by

$$I_f(x) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} f_{mnk} (|x_{mnk}|)^{1/m+n+k}, x = (x_{mnk}) \in t_f.$$

We consider t_f equipped with the Luxemburg metric

$$d(x, y) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} f_{mnk} \left(\frac{|x_{mnk}|^{1/m+n+k}}{mnk} \right)$$

is an extended real number.

Definition 2.2. Let α be a proper fraction. A triple difference sequence spaces of $\Delta_\gamma^\alpha x = (\Delta_\gamma^\alpha x_{mnk})$ is said to be Δ_γ^α strong Cesàro summable to $\bar{0}$ if

$$\lim_{uvw \rightarrow \infty} \frac{1}{uvw} \sum_{m=1}^u \sum_{n=1}^v \sum_{k=1}^w |\Delta_\gamma^\alpha x_{mnk}, \bar{0}| = 0.$$

In this we write $\Delta_\gamma^\alpha x_{mnk} \xrightarrow{[C,1,1,1]} \Delta_\gamma^\alpha x_{mnk}$. The set of all Δ_γ^α strong Cesàro summable triple sequence spaces is denoted by $[C, 1, 1, 1]$.

Definition 2.3. Let α be a proper fraction and β be a nonnegative real number. A triple difference sequence spaces of $\Delta_\gamma^\alpha x = (\Delta_\gamma^\alpha x_{mnk})$ is said to be Δ_γ^α rough strong Cesàro summable in probability to a random variable $\Delta_\gamma^\alpha x : W \times W \times W \rightarrow \mathbb{R} \times \mathbb{R} \times \mathbb{R}$ with respect to the roughness of degree β if for each $\epsilon > 0$,

$$\lim_{uvw \rightarrow \infty} \frac{1}{uvw} \sum_{m=1}^u \sum_{n=1}^v \sum_{k=1}^w P(|\Delta_\gamma^\alpha x_{mnk}, \bar{0}| \geq \beta + \epsilon) = 0.$$

In this case we write $\Delta_\gamma^\alpha x_{mnk} \xrightarrow{\beta [C,1,1,1]^{P\Delta}} \Delta_\gamma^\alpha x_{mnk}$. The class of all $\beta \Delta_\gamma^\alpha$ -strong Cesàro summable triple sequence spaces of random variables in probability and it will be denoted by $\beta [C, 1, 1, 1]^{P\Delta}$.

3. Rough Cesàro Summable of Triple of Δ_γ^α

In this section by using the operator Δ_γ^α , we introduce some new triple difference sequence spaces of rough Cesàro summable involving lacunary sequences θ and arbitrary sequence $p = (p_{rst})$ of strictly positive real numbers.

If α be a proper fraction and β be nonnegative real number. A triple difference sequence spaces of $\Delta_\gamma^\alpha X = (\Delta_\gamma^\alpha x_{mnk})$ is said to be Δ_γ^α -rough strong Cesàro summable in probability to a random variable $\Delta_\gamma^\alpha X : W \times W \times W \rightarrow \mathbb{R} \times \mathbb{R} \times \mathbb{R}$ with respect to the roughness of degree β if for each $\epsilon > 0$ then define the triple difference sequence spaces as follows:

(i)

$$C(\Delta_\gamma^\alpha, p)_\theta = \sum_{r=1}^{\infty} \sum_{s=1}^{\infty} \sum_{t=1}^{\infty} P \left(f_{mnk} \left[\left| \frac{1}{h_{rst}} \sum_{(mnk) \in I_{rst}} \Delta_\gamma^\alpha X \right|^{p_{rst}} \right] \geq \beta + \epsilon \right) < \infty.$$

In this case we write $C(\Delta_\gamma^\alpha, p)_\theta \xrightarrow{[C,1,1,1]^{P\Delta}} C(\Delta_\gamma^\alpha, p)_\theta$. The class of all $\beta C(\Delta_\gamma^\alpha, p)_\theta$ – rough strong Cesàro summable triple sequence spaces of random variables in probability and it will be denoted by $\beta[C, 1, 1, 1]^{P\Delta}$.

(ii)

$$C[\Delta_\gamma^\alpha, p]_\theta = \sum_{r=1}^{\infty} \sum_{s=1}^{\infty} \sum_{t=1}^{\infty} P \left(\frac{1}{h_{rst}} \sum_{(mnk) \in I_{rst}} f_{mnk} [|\Delta_\gamma^\alpha X|^{p_{rst}}] \geq \beta + \epsilon \right) < \infty.$$

In this case we write $C[\Delta_\gamma^\alpha, p]_\theta \xrightarrow{[C,1,1,1]^{P\Delta}} C[\Delta_\gamma^\alpha, p]_\theta$. The class of all $\beta C[\Delta_\gamma^\alpha, p]_\theta$ – rough strong Cesàro summable triple sequence spaces of random variables in probability.

(iii)

$$C_\Lambda(\Delta_\gamma^\alpha, p)_\theta = P \left(f_{mnk} \left[\left| \frac{1}{h_{rst}} \sum_{(mnk) \in I_{rst}} \Delta_\gamma^\alpha X \right|^{p_{rst}} \right] \geq \beta + \epsilon \right) < \infty.$$

In this case we write $C_\Lambda(\Delta_\gamma^\alpha, p)_\theta \xrightarrow{[C,1,1,1]^{P\Delta}} C_\Lambda(\Delta_\gamma^\alpha, p)_\theta$. The class of all $\beta C_\Lambda(\Delta_\gamma^\alpha, p)_\theta$ – rough strong Cesàro summable triple sequence spaces of random variables in probability.

(iv)

$$C_\Lambda[\Delta_\gamma^\alpha, p]_\theta = \frac{1}{h_{rst}} \sum_{(mnk) \in I_{rst}} P(f_{mnk} [|\Delta_\gamma^\alpha X|^{p_{rst}}] \geq \beta + \epsilon) < \infty.$$

In this case we write $C_\Lambda[\Delta_\gamma^\alpha, p]_\theta \xrightarrow{[C,1,1,1]^{P\Delta}} C_\Lambda[\Delta_\gamma^\alpha, p]_\theta$. The class of all $\beta C_\Lambda[\Delta_\gamma^\alpha, p]_\theta$ – rough strong Cesàro summable triple sequence spaces of random variables in probability.

(v)

$$N(\Delta_\gamma^\alpha, p)_\theta = \lim_{rst \rightarrow \infty} \frac{1}{h_{rst}} \sum_{(mnk) \in I_{rst}} P(f_{mnk} [|\Delta_\gamma^\alpha X, \bar{0}|^{p_{rst}}] \geq \beta + \epsilon) = 0.$$

In this case we write $N(\Delta_\gamma^\alpha, p)_\theta \xrightarrow{[C,1,1,1]^{P\Delta}} N(\Delta_\gamma^\alpha, p)_\theta$. The class of all $\beta N(\Delta_\gamma^\alpha, p)_\theta$ – rough strong Cesàro summable triple sequence spaces of random variables in probability.

Theorem 3.1. *If α be a proper fraction, β be nonnegative real number, f be an Musielak-Orlicz function and (p_{rst}) is a triple difference analytic sequence then the sequence spaces $C(\Delta_\gamma^\alpha, p)_\theta$, $C[\Delta_\gamma^\alpha, p]_\theta$, $C_\Lambda(\Delta_\gamma^\alpha, p)_\theta$, $C_\Lambda[\Delta_\gamma^\alpha, p]_\theta$ and $N(\Delta_\gamma^\alpha, p)_\theta$ are linear spaces.*

Proof: Because the linearity may be proved in a similar way for each of the sets of triple sequences, hence it is omitted.

Theorem 3.2. *If α be a proper fraction, β be nonnegative real number, f be an Musielak-Orlicz function and (p_{rst}) , for all $r, s, t \in \mathbb{N}$, then the triple difference sequence spaces $C[\Delta_\gamma^\alpha, p]_\theta$ is a BK-space with the Luxemburg metric is defined by*

$$d(x, y)_1 = \sum_{u=0}^{\infty} \sum_{v=0}^{\infty} \sum_{w=0}^{\infty} f_{mnk} \left[\frac{\gamma_{uvw} x_{uvw}}{uvw} \right] \\ + \lim_{uvw \rightarrow \infty} \frac{1}{uvw} \sum_{r=1}^{\infty} \sum_{s=1}^{\infty} \sum_{t=1}^{\infty} f_{mnk} \left[P \left(\frac{1}{h_{rst}} \sum_{(m,n,k) \in I_{rst}} |\Delta_\gamma^\alpha x|^p \right) \geq \beta + \epsilon \right]^{1/p}, \quad 1 \leq p.$$

Also if $p_{rst} = 1$ for all $(r, s, t) \in \mathbb{N}$, then the triple difference spaces $C_\Lambda[\Delta_\gamma^\alpha, p]_\theta$ and $N(\Delta_\gamma^\alpha, p)_\theta$ are BK-spaces with the Luxemburg metric is defined by

$$d(x, y)_2 = \sum_{u=0}^{\infty} \sum_{v=0}^{\infty} \sum_{w=0}^{\infty} f_{mnk} \left[\frac{\gamma_{uvw} x_{uvw}}{uvw} \right] \\ + \lim_{uvw \rightarrow \infty} \frac{1}{uvw} \frac{1}{h_{rst}} \sum_{(m,n,k) \in I_{rst}} f_{mnk} [P(|\Delta_\gamma^\alpha x|) \geq \beta + \epsilon].$$

Proof. We give the proof for the space $C_\Lambda[\Delta_\gamma^\alpha, p]_\theta$ and that of others followed by using similar techniques.

Suppose (x^n) is a Cauchy sequence in $C_\Lambda[\Delta_\gamma^\alpha, p]_\theta$, where $x^n = (x_{ij\ell})^n$ and $x^m = (x_{ij\ell}^m)$ are two elements in $C_\Lambda[\Delta_\gamma^\alpha, p]_\theta$. Then there exists a positive integer $n_0(\epsilon)$ such that $|x^n - x^m| \rightarrow 0$ as $m, n \rightarrow \infty$ for all $m, n \geq n_0(\epsilon)$ and for each $i, j, \ell \in \mathbb{N}$. Therefore

$$\begin{bmatrix} x_{uvw}^{11} & x_{uvw}^{12} & \dots & \dots \\ x_{uvw}^{21} & x_{uvw}^{22} & \dots & \dots \\ \vdots & \vdots & \ddots & \ddots \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} \Delta_\gamma^\alpha x_{ij\ell}^{11} & \Delta_\gamma^\alpha x_{ij\ell}^{12} & \dots & \dots \\ \Delta_\gamma^\alpha x_{ij\ell}^{21} & \Delta_\gamma^\alpha x_{ij\ell}^{22} & \dots & \dots \\ \vdots & \vdots & \ddots & \ddots \end{bmatrix}$$

are Cauchy sequences in complex field \mathbb{C} and $C_\Lambda[\Delta_\gamma^\alpha, p]_\theta$ respectively. By using the completeness of \mathbb{C} and $C_\Lambda[\Delta_\gamma^\alpha, p]_\theta$ we have that they are convergent and suppose

that $x_{ij\ell}^n \rightarrow x_{ij\ell}$ in \mathbb{C} and $(\Delta_\gamma^\alpha x_{ij\ell}^n) \rightarrow y_{ij\ell}$ in $C_\Lambda [\Delta_\gamma^\alpha, p]_\theta$ for each $i, j, \ell \in \mathbb{N}$ as $n \rightarrow \infty$. Then we can find a triple sequence space of $(x_{ij\ell})$ such that $y_{ij\ell} = \Delta_\gamma^\alpha x_{ij\ell}$ for $i, j, \ell \in \mathbb{N}$. These $x_{ij\ell}^n$ can be interpreted as

$$x_{ij\ell} = \frac{1}{\gamma_{ij\ell}} \sum_{u=1}^{i-m} \sum_{v=1}^{j-n} \sum_{w=1}^{\ell-k} \Delta_\gamma^\alpha y_{uvw}$$

$$= \frac{1}{\gamma_{ij\ell}} \sum_{u=1}^i \sum_{v=1}^j \sum_{w=1}^\ell \Delta_\gamma^\alpha y_{u-m, v-n, w-k}, (y_{1-m, 1-n, 1-k} = y_{2-m, 2-n, 2-k} = \dots = y_{000} = 0).$$

for sufficiently large (i, j, ℓ) ; that is,

$$(\Delta_\gamma^\alpha x^n) = \begin{bmatrix} \Delta_\gamma^\alpha x_{ij\ell}^{11} & \Delta_\gamma^\alpha x_{ij\ell}^{12} & \dots & \dots \\ \Delta_\gamma^\alpha x_{ij\ell}^{21} & \Delta_\gamma^\alpha x_{ij\ell}^{22} & \dots & \dots \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix}$$

converges to $(\Delta_\gamma^\alpha x_{ij\ell})$ for each $i, j, \ell \in \mathbb{N}$ as $n \rightarrow \infty$. Thus $|x^m - x|_2 \rightarrow 0$ as $m \rightarrow \infty$. Since $C_\Lambda [\Delta_\gamma^\alpha, p]_\theta$ is a Banach Luxemburg metric with continuous coordinates, that is $|x^n - x|_2 \rightarrow 0$ implies $|x_{ij\ell}^n - x_{ij\ell}| \rightarrow 0$ for each $i, j, \ell \in \mathbb{N}$ as $n \rightarrow \infty$, this shows that $C_\Lambda [\Delta_\gamma^\alpha, p]_\theta$ is a BK-space. \square

Theorem 3.3. *If α be a proper fraction, β be nonnegative real number, f be an Musielak-Orlicz function and (p_{rst}) , for all $r, s, t \in \mathbb{N}$, then the triple difference sequence space $C(\Delta_\gamma^\alpha, p)_\theta$ is a BK-space with the Luxemburg metric is defined by*

$$d(x, y)_3 = \sum_{u=0}^\infty \sum_{v=0}^\infty \sum_{w=0}^\infty f_{mnk} \left[\frac{\gamma_{uvw} x_{uvw}}{uvw} \right]$$

$$+ \lim_{uvw \rightarrow \infty} \frac{1}{uvw} \sum_{r=1}^\infty \sum_{s=1}^\infty \sum_{t=1}^\infty f_{mnk} \left[P \left(\left| \frac{1}{h_{rst}} \sum_{(m,n,k) \in I_{rst}} \Delta_\gamma^\alpha x \right|^p \right) \geq \beta + \epsilon \right]^{1/p}, \quad 1 \leq p.$$

Also if $p_{rst} = 1$ for all $(r, s, t) \in \mathbb{N}$, then the triple difference spaces $C_\Lambda (\Delta_\gamma^\alpha, p)_\theta$ is a BK-spaces with the Luxemburg metric is defined by

$$d(x, y)_4 = \sum_{u=0}^\infty \sum_{v=0}^\infty \sum_{w=0}^\infty f_{mnk} \left[\frac{\gamma_{uvw} X_{uvw}}{uvw} \right]$$

$$+ \lim_{uvw \rightarrow \infty} \frac{1}{uvw} f_{mnk} \left[P \left(\left| \frac{1}{h_{rst}} \sum_{(m,n,k) \in I_{rst}} \Delta_\gamma^\alpha x \right| \right) \geq \beta + \epsilon \right].$$

Proof: The proof follows from Theorem 4.2.

Now, we can present the following theorem, determining some inclusion relations without proof, since it is a routine verification.

Theorem 3.4. *Let α, ξ be two positive proper fractions $\alpha > \xi > 0$ and β be two nonnegative real number, f be an Musielak-Orlicz function and $(p_{rst}) = p$, for each $r, s, t \in \mathbb{N}$ be given. Then the following inclusions are satisfied:*

- (i) $C(\Delta_\gamma^\xi, p)_\theta \subset C(\Delta_\gamma^\alpha, p)_\theta$;
- (ii) $C[\Delta_\gamma^\xi, p]_\theta \subset C[\Delta_\gamma^\alpha, p]_\theta$;
- (iii) $C(\Delta_\gamma^\alpha, p)_\theta \subset C(\Delta_\gamma^\alpha, q)_\theta$, $0 < p < q$.

4. Rough Lacunary Statistical Convergence of Triple of Δ_γ^α

In this section by using the operator Δ_γ^α , we introduce some new triple difference sequence spaces involving rough lacunary statistical sequences spaces and arbitrary sequence $p = (p_{rst})$ of strictly positive real numbers.

Definition 4.1. The triple sequence $\theta_{i,\ell,j} = \{(m_i, n_\ell, k_j)\}$ is called triple lacunary if there exist three increasing sequences of integers such that

$$\begin{aligned} m_0 &= 0, h_i = m_i - m_{r-1} \rightarrow \infty \text{ as } i \rightarrow \infty \text{ and} \\ n_0 &= 0, \bar{h}_\ell = n_\ell - n_{\ell-1} \rightarrow \infty \text{ as } \ell \rightarrow \infty, \\ k_0 &= 0, \bar{h}_j = k_j - k_{j-1} \rightarrow \infty \text{ as } j \rightarrow \infty. \end{aligned}$$

Let $m_{i,\ell,j} = m_i n_\ell k_j$, $h_{i,\ell,j} = h_i \bar{h}_\ell \bar{h}_j$, and $\theta_{i,\ell,j}$ is determine by

$$\begin{aligned} I_{i,\ell,j} &= \{(m, n, k) : m_{i-1} < m < m_i \text{ and } n_{\ell-1} < n \leq n_\ell \text{ and } k_{j-1} < k \leq k_j\}, \\ q_i &= \frac{m_i}{m_{i-1}}, \bar{q}_\ell = \frac{n_\ell}{n_{\ell-1}}, \bar{q}_j = \frac{k_j}{k_{j-1}}. \end{aligned}$$

Definition 4.2. Let α be a proper fraction, f be an Musielak-Orlicz function and $\theta = \{m_r n_s k_t\}_{(rst) \in \mathbb{N} \cup 0}$ be the triple difference lacunary sequence spaces of $(\Delta_\gamma^\alpha X_{mnk})$ is said to be Δ_γ^α -lacunary statistically convergent to a number $\bar{0}$ if for any $\epsilon > 0$,

$$\lim_{rst \rightarrow \infty} \frac{1}{h_{rst}} |\{(m, n, k) \in I_{rst} : f_{mnk} [|\Delta_\gamma^\alpha X_{mnk}, \bar{0}|] \geq \epsilon\}| = 0,$$

where

$$\begin{aligned} I_{r,s,t} &= \{(m, n, k) : m_{r-1} < m < m_r \text{ and } n_{s-1} < n \leq n_s \text{ and } k_{t-1} < k \leq k_t\}, \\ q_r &= \frac{m_r}{m_{r-1}}, \bar{q}_s = \frac{n_s}{n_{s-1}}, \bar{q}_t = \frac{k_t}{k_{t-1}}. \end{aligned}$$

In this case write $\Delta_\gamma^\alpha X \rightarrow^{S_\theta} \Delta_\gamma^\alpha x$.

Definition 4.3. If α be a proper fraction, β be nonnegative real number, f be an Musielak-Orlicz function and $\theta = \{m_r n_s k_t\}_{(r,s,t) \in \mathbb{N}^3 \cup (0,0,0)}$ be the triple difference sequence spaces of lacunary. A number X is said to be $\Delta_\gamma^\alpha - N_\theta$ -convergent to a real number $\bar{0}$ if for every $\epsilon > 0$,

$$\lim_{rst \rightarrow \infty} \frac{1}{h_{rst}} \sum_{m \in I_r} \sum_{n \in I_s} \sum_{k \in I_t} f_{mnk} [|\Delta_\gamma^\alpha X_{mnk}, \bar{0}|] = 0.$$

In this case we write $\Delta_\gamma^\alpha X_{mnk} \rightarrow^{N_\theta} \bar{0}$.

Definition 4.4. Let α be a proper fraction, β be nonnegative real number, f be an Musielak-Orlicz function and arbitrary sequence $p = (p_{rst})$ of strictly positive real numbers. A triple difference sequence spaces of random variables is said to be Δ_γ^α -rough lacunary statistically convergent in probability to $\Delta_\gamma^\alpha X : W \times W \times W \rightarrow \mathbb{R} \times \mathbb{R} \times \mathbb{R}$ with respect to the roughness of degree β if for any $\epsilon, \delta > 0$,

$$\lim_{rst \rightarrow \infty} \frac{1}{h_{rst}} |\{(m, n, k) \in I_{rst} : P([f_{mnk} (|\Delta_\gamma^\alpha(x_{mnk})|)]^{p_{rst}} \geq \beta + \epsilon) \geq \delta\}| = 0$$

and we write $\Delta_\gamma^\alpha X_{mnk} \rightarrow_{\beta}^{S^P} \bar{0}$. It will be denoted by βS_θ^P .

Definition 4.5. Let α be a proper fraction, β be nonnegative real number, f be an Musielak-Orlicz function and arbitrary sequence $p = (p_{rst})$ of strictly positive real numbers. A triple difference sequence spaces of random variables is said to be Δ_γ^α -rough N_θ -convergent in probability to $\Delta_\gamma^\alpha X : W \times W \times W \rightarrow \mathbb{R} \times \mathbb{R} \times \mathbb{R}$ with respect to the roughness of degree β if for any $\epsilon > 0$,

$$\lim_{rst \rightarrow \infty} \frac{1}{h_{rst}} \sum_{m \in I_r} \sum_{n \in I_s} \sum_{k \in I_t} |\{P([f_{mnk} (|\Delta_\gamma^\alpha X_{mnk}|)]^{p_{rst}} \geq \beta + \epsilon)\}| = 0,$$

and we write $\Delta_\gamma^\alpha X_{mnk} \rightarrow_{\beta}^{N_\theta^P} \Delta_\gamma^\alpha X$. The class of all $\beta - N_\theta$ -convergent triple difference sequence spaces of random variables in probability will be denoted by βN_θ^P .

Definition 4.6. Let α be a proper fraction, β be nonnegative real number, f be an Musielak-Orlicz function and arbitrary sequence $p = (p_{rst})$ of strictly positive real numbers. A triple difference sequence spaces of random variables is said to be Δ_γ^α -rough lacunary statistically Cauchy if there exists a number $N = N(\epsilon)$ in probability to $\Delta_\gamma^\alpha X : W \times W \times W \rightarrow \mathbb{R} \times \mathbb{R} \times \mathbb{R}$ with respect to the roughness of degree β if for any $\epsilon, \delta > 0$,

$$\lim_{rst \rightarrow \infty} \frac{1}{h_{rst}} |\{(m, n, k) \in I_{rst} : P([f_{mnk} (|\Delta_\gamma^\alpha(x_{mnk} - x_N)|)]^{p_{rst}} \geq \beta + \epsilon) \geq \delta\}| = 0.$$

Theorem 4.1. Let α be a proper fraction, β be nonnegative real number, f be an Musielak-Orlicz function and arbitrary sequence $p = (p_{rst})$ of strictly positive real numbers, $0 < p < \infty$.

(i) If $(x_{mnk}) \rightarrow (N(\Delta_\gamma^\alpha, p)_\theta)$ for $p_{rst} = p$ then $(x_{mnk}) \rightarrow (\Delta_\gamma^\alpha(S_\theta))$.

(ii) If $x \in (\Delta_\gamma^\alpha(S_\theta))$, then $(x_{mnk}) \rightarrow (N(\Delta_\gamma^\alpha, p)_\theta)$.

Proof. Let $x = (x_{mnk}) \in (N(\Delta_\gamma^\alpha, p)_\theta)$ and $\epsilon > 0$,

$$|\{P([f_{mnk}(|\Delta_\gamma^\alpha X_{mnk}|)]^{p_{rst}} \geq \beta + \epsilon)\}| = 0.$$

We have

$$\begin{aligned} & \frac{1}{h_{rst}} \sum_{(mnk) \in I_{rst}} |\{P([f_{mnk}(|\Delta_\gamma^\alpha X_{mnk}|)]^{p_{rst}} \geq \beta + \epsilon)\}| \\ & \geq \frac{1}{h_{rst}} |\{(m, n, k) \in I_{rst} : P([f_{mnk}(|\Delta_\gamma^\alpha(x_{mnk})|)]^{p_{rst}} \geq \beta + \epsilon) \geq \delta\}| \left(\frac{\beta + \epsilon}{\delta}\right)^p. \end{aligned}$$

So we observe by passing to limit as $r, s, t \rightarrow \infty$,

$$\begin{aligned} & \lim_{rst \rightarrow \infty} \frac{1}{h_{rst}} |\{(m, n, k) \in I_{rst} : P([f_{mnk}(|\Delta_\gamma^\alpha(x_{mnk})|)]^{p_{rst}} \geq \beta + \epsilon) \geq \delta\}| \\ & \leq \left(\frac{\delta}{\alpha + \epsilon}\right)^p P\left(\lim_{rst \rightarrow \infty} \frac{1}{h_{rst}} \sum_{(m, n, k) \in I_{rst}} |\Delta_\gamma^\alpha x_{mnk}|^p\right) = 0, \end{aligned}$$

which implies that $x_{mnk} \rightarrow (\Delta_\gamma^\alpha(S_\theta))$.

Suppose that $x \in \Delta_\gamma^\alpha(\Lambda^3)$ and $(x_{mnk}) \rightarrow (\Delta_\gamma^\alpha(S))$. Then it is obvious that $(\Delta_\gamma^\alpha x) \in \Lambda^3$ and

$$\frac{1}{h_{rst}} |\{(m, n, k) \in I_{rst} : P([f_{mnk}(|\Delta_\gamma^\alpha(x_{mnk})|)]^{p_{rst}} \geq \beta + \epsilon) \geq \delta\}| \rightarrow 0$$

as $r, s, t \rightarrow \infty$. Let $\epsilon > 0$ be given and there exists $u_0 v_0 w_0 \in \mathbb{N}$ such that

$$\begin{aligned} & \left| \left\{ (m, n, k) \in I_{rst} : P\left([f_{mnk}(|\Delta_\gamma^\alpha(x_{mnk})|)]^{p_{rst}} \geq \beta + \frac{\epsilon}{2}\right) \geq \frac{\delta}{2} \right\} \right| \\ & \leq \frac{\epsilon}{2(d(\Delta_\gamma^\alpha x, y))_{\Lambda^3}} + \frac{\delta}{2}, \end{aligned}$$

where $\sum_{u=1}^\infty \sum_{v=1}^\infty \sum_{w=1}^\infty |\gamma_{uvw} x_{uvw}| = 0$, for all $r \geq u_0, s \geq v_0, t \geq w_0$. Further more, we can write $|\Delta_\gamma^\alpha x_{mnk}| \leq d(\Delta_\gamma^\alpha x_{mnk}, y)_{\Delta_\gamma^\alpha} \leq d(\Delta_\gamma^\alpha x, y)_{\Lambda^3} = d(x, y)_{\Delta_\gamma^\alpha x}$. For $r, s, t \geq u_0, v_0, w_0$

$$\begin{aligned} & \frac{1}{h_{rst}} \sum_{(mnk) \in I_{rst}} P([f_{mnk}(|\Delta_\gamma^\alpha X_{mnk}|)]^p) = \frac{1}{h_{rst}} P\left(\sum_{(mnk) \in I_{rst}} [f_{mnk}(|\Delta_\gamma^\alpha X_{mnk}|)]^p\right) \\ & + \frac{1}{h_{rst}} P\left(\sum_{(mnk) \notin I_{rst}} [f_{mnk}(|\Delta_\gamma^\alpha X_{mnk}|)]^p\right) < \frac{1}{h_{rst}} P\left(h_{rst} \left(\frac{\epsilon}{2} + \frac{\delta}{2}\right)\right) \\ & + h_{rst} \frac{\epsilon d(x, y)_{\Delta_\gamma^\alpha X}^p}{2 d(x, y)_{\Delta_\gamma^\alpha X}^p} + \frac{\delta}{2} = \epsilon + \delta. \end{aligned}$$

Hence $(x_{mnk}) \rightarrow (N(\Delta_\gamma^\alpha, p)_\theta)$. □

Corollary 4.1. *If α be a proper fraction, β be nonnegative real number, f be an Musielak-Orlicz function and arbitrary sequence $p = (p_{rst})$ of strictly positive real numbers then the following statements are hold:*

- (i) $S \cap \Lambda^3 \subset \Delta_\gamma^\alpha(S_\theta) \cap \Delta_\gamma^\alpha(\Lambda^3)$;
- (ii) $\Delta_\gamma^\alpha(S_\theta) \cap \Delta_\gamma^\alpha(\Lambda^3) = \Delta_\gamma^\alpha(w_p^3)$.

Theorem 4.2. *Let α be a proper fraction, β be nonnegative real number, f be an Musielak-Orlicz function and arbitrary sequence $p = (p_{rst})$ of strictly positive real numbers. If $x = (x_{mnk})$ is a Δ_γ^α -triple difference rough lacunary statistically convergent sequence, then x is a Δ_γ^α -triple difference rough lacunary statistically Cauchy sequence.*

Proof. Assume that $(x_{mnk}) \rightarrow (\Delta_\gamma^\alpha(S_\theta))$ and $\epsilon, \delta > 0$. Then

$$\frac{1}{\delta} \left| \left\{ (m, n, k) \in I_{rst} : P \left([f_{mnk} (|\Delta_\gamma^\alpha x_{mnk}|)]^{p_{rst}} \geq \beta + \frac{\epsilon}{2} \right) \right\} \right|$$

for almost all m, n, k and if we select N , then

$$\frac{1}{\delta} \left| \left\{ (m, n, k) \in I_{rst} : P \left([f_{mnk} (|\Delta_\gamma^\alpha x_N|)]^{p_{rst}} \geq \beta + \frac{\epsilon}{2} \right) \right\} \right|$$

holds. Now, we have

$$\begin{aligned} & \left| \left\{ (m, n, k) \in I_{rst} : P \left([f_{mnk} (|\Delta_\gamma^\alpha (x_{mnk} - x_N)|)]^{p_{rst}} \right) \right\} \right| \\ & \leq \frac{1}{\delta} \left| \left\{ (m, n, k) \in I_{rst} : P \left([f_{mnk} (|\Delta_\gamma^\alpha x_{mnk}|)]^{p_{rst}} \geq \beta + \frac{\epsilon}{2} \right) \right\} \right| \\ & + \frac{1}{\delta} \left| \left\{ (m, n, k) \in I_{rst} : P \left([f_{mnk} (|\Delta_\gamma^\alpha x_N|)]^{p_{rst}} \geq \beta + \frac{\epsilon}{2} \right) \right\} \right| < \frac{1}{\delta} (\beta + \epsilon) = \epsilon, \end{aligned}$$

for almost all m, n, k . Hence (x_{mnk}) is a Δ_γ^α -rough lacunary statistically Cauchy. □

Theorem 4.3. *If α be a proper fraction, β be nonnegative real number, f be an Musielak-Orlicz function and arbitrary sequence $p = (p_{rst})$ of strictly positive real numbers and $0 < p < \infty$, then $N(\Delta_\gamma^\alpha, p)_\theta \subset \Delta_\gamma^\alpha(S_\theta)$.*

Proof. Suppose that $x = (x_{mnk}) \in N(\Delta_\gamma^\alpha, p)_\theta$ and

$$\left| \left\{ (m, n, k) \in I_{rst} : P \left([f_{mnk} (|\Delta_\gamma^\alpha x_{mnk}|)]^p \geq \beta + \epsilon \right) \right\} \right|.$$

Therefore we have

$$\begin{aligned} & \frac{1}{h_{rst}} \sum_{(mnk) \in I_{rst}} P \left([f_{mnk} (|\Delta_\gamma^\alpha x_{mnk}|)]^p \right) \geq \frac{1}{h_{rst}} \sum_{(mnk) \in I_{rst}} (\beta + \epsilon)^p \\ & \geq \frac{1}{h_{rst}} \left| \left\{ (m, n, k) \in I_{rst} : P \left([f_{mnk} (|\Delta_\gamma^\alpha x_{mnk}|)]^p \geq \beta + \epsilon \right) \right\} \right| (\beta + \epsilon)^p. \end{aligned}$$

So we observe by passing to limit as $r, s, t \rightarrow \infty$,

$$\begin{aligned} & \lim_{rst \rightarrow \infty} \frac{1}{h_{rst}} |\{(m, n, k) \in I_{rst} : P([f_{mnk}(|\Delta_\gamma^\alpha(x_{mnk})|)]^p \geq \beta + \epsilon) \geq \delta\}| \\ & < \frac{1}{(\beta + \epsilon)^p} \left(P \left(\lim_{rst \rightarrow \infty} \frac{1}{h_{rst}} \sum_{(m,n,k) \in I_{rst}} [f_{mnk}(|\Delta_\gamma^\alpha(x_{mnk})|)]^p \right) \right) = 0 \end{aligned}$$

implies that $x \in \Delta_\gamma^\alpha(S_\theta)$. Hence $N(\Delta_\gamma^\alpha, p)_\theta \subset \Delta_\gamma^\alpha(S_\theta)$. \square

Competing Interests: The authors declare that there is not any conflict of interests regarding the publication of this manuscript.

References

- [1] A. Esi, *On some triple almost lacunary sequence spaces defined by Orlicz functions*, Research and Reviews: Discrete Mathematical Structures 1 (2) (2014) 16–25.
- [2] A. Esi, M. Necdet Catalbas, *Almost convergence of triple sequences*, Global Journal of Mathematical Analysis 2 (1) (2014) 6–10.
- [3] A. Esi, E. Savas, *On lacunary statistically convergent triple sequences in probabilistic normed space*, Appl. Math. and Inf. Sci. 9 (5) (2015) 2529–2534.
- [4] A.J. Dutta, A. Esi, B.C. Tripathy, *Statistically convergent triple sequence spaces defined by Orlicz function*, Journal of Mathematical Analysis 4 (2) (2013) 16–22.
- [5] S. Debnath, B. Sarma, B.C. Das, *Some generalized triple sequence spaces of real numbers*, Journal of Nonlinear Analysis and Optimization 6 (1) (2015) 71–79.
- [6] H. Kizmaz, *On certain sequence spaces*, Canadian Mathematical Bulletin 24 (2) (1981) 169–176.
- [7] P.K. Kamthan, M. Gupta, *Sequence Spaces and Series*, Lecture Notes, Pure and Applied Mathematics, 65 Marcel Dekker Inc. New York, 1981.
- [8] J. Lindenstrauss, L. Tzafriri, *On Orlicz sequence spaces*, Israel J. Math. 10 (1971) 379–390.
- [9] J. Musielak, *Orlicz Spaces*, Lectures Notes in Math., 1034, Springer-Verlag, 1983.
- [10] A. Sahiner, M. Gurdal, F.K. Duden, *Triple sequences and their statistical convergence*, Selcuk J. Appl. Math. 8 (2) (2007) 49–55.
- [11] A. Sahiner, B.C. Tripathy, *Some I-related properties of triple sequences*, Selcuk J. Appl. Math. 9 No. (2) (2008) 9–18.

- [12] N. Subramanian, A. Esi, *The generalized tripled difference of χ^3 sequence spaces*, Global Journal of Mathematical Analysis 3 (2) (2015) 54–60.
- [13] A. Esi, N. Subramanian, *The triple sequence spaces of χ^3 on rough statistical convergence defined by Musielak Orlicz function of p -metric*, Asian Journal of Mathematical Sciences 1 (1) (2017) 19–25.
- [14] N. Subramanian, A. Esi, M.K. Ozdemir, *Some new triple intuitionistic sequence spaces of fuzzy numbers defined by Musielak-Orlicz function*, J. Assam Acad. Math. 7 (2017) 14–27.
- [15] A. Esi, N. Subramanian, A. Esi, *Triple rough statistical convergence of sequence of Bernstein operators*, Int. J. Adv. Appl. Sci. 4 (2) (2017) 28–34.

DOI: 10.7862/rf.2018.7

Ayhan Esi

email: aesi23@hotmail.com
ORCID: 0000-0003-3137-3865
Department of Mathematics
Adiyaman University
02040, Adiyaman
TURKEY

Nagarajan Subramanian

email: nsmaths@yahoo.com
ORCID: 0000-0002-5895-673X
Department of Mathematics
SASTRA University
Thanjavur-613 401
INDIA

Received 02.07.2018

Accepted 21.10.2018